

Fermions Tunneling from Apparent Horizon of FRW Universe

Ran Li, Ji-Rong Ren,^{*} and Dun-Fu Shi

Institute of Theoretical Physics, Lanzhou University, Lanzhou, 730000, Gansu, China

Abstract

In the paper [arXiv:0809.1554], the scalar particles' Hawking radiation from the apparent horizon of Friedmann-Robertson-Walker(FRW) universe was investigated by using the tunneling formalism. They obtained the Hawking temperature associated with the apparent horizon, which was extensively applied in investigating the relationship between the first law of thermodynamics and Friedmann equations. In this paper, we calculate Fermions' Hawking radiation from the apparent horizon of FRW universe via tunneling formalism. Applying WKB approximation to the general covariant Dirac equation in FRW spacetime background, the radiation spectrum and Hawking temperature of apparent horizon are correctly recovered, which supports the arguments presented in the paper [arXiv:0809.1554].

PACS numbers: 04.62.+v, 04.70.Dy, 98.80.Jk.

Keywords: Fermions, tunneling, Hawking radiation, FRW Universe.

^{*}Corresponding author. Electronic mail: renjr@lzu.edu.cn

In the recent paper [1], Cai *et al* discussed the scalar particles' Hawking radiation from the apparent horizon of FRW universe, which revealed that there is a Hawking temperature associated with the apparent horizon. In this paper, we will investigate Fermions' tunneling from the apparent horizon of FRW universe. The results in the present paper together with that in [1] fill in the gap existing in the literatures investigating the relationship between the first law of thermodynamics and Friedmann equations of FRW universe.

Ted Jacobson[2] has been able to derive Einstein equation from the proportionality of entropy and horizon area together with the fundamental Clausius relation $\delta Q = TdS$ connecting heat, entropy, and temperature. The key idea is to demand that this relation holds for all the local Rindler causal horizons through each spacetime point, with δQ and T interpreted as the energy flux and Unruh temperature seen by an accelerated observer just inside the horizon. This perspective suggests that Einstein field equation can be viewed as an equation of state of spacetime. The idea has also applied to $f(R)$ theory[3] and scalar-tensor theory[4], where the non-equilibrium thermodynamic must be taken into account.

It has been proved that the above idea can also be applied in establishing the relationship between the Friedmann equations and the first law of thermodynamics in the framework of Friedmann-Robertson-Walker(FRW) universe. Assuming that the apparent horizon of FRW universe has the temperature $T = 1/2\pi\tilde{r}_A$ and entropy $S = A/4$, where $\tilde{r}_A$ and A are the radius and the area of the apparent horizon respectively, the Friedmann equations can be derived from the Clausius relation[5] and the first Friedmann equation can be cast into the form of the unified first law[4, 6]. For some related investigations see also[7, 8, 9, 10, 11, 12, 13, 14, 15].

However, whether there is a Hawking temperature associated with the apparent horizon of FRW universe is still an open question to investigate. In a recent paper [1], the scalar particles' Hawking radiation from the Apparent Horizon of FRW Universe was investigated by using the tunneling formalism. They obtained the Hawking temperature associated with the apparent horizon of FRW universe. They found that the Kodama observer inside the apparent horizon do see a thermal spectrum with temperature $T = 1/2\pi\tilde{r}_A$, which is caused by particles tunneling from the outside apparent horizon to the inside apparent horizon.

The semi-classical derivation of Hawking radiation as tunneling process[16] was initially proposed by Parikh and Wilczek. In recent years, it has already attracted a lot of attention. In this method, the imaginary part of the action is calculated using the null geodesic

equation. Zhang and Zhao extended this method to the charged Reissner-Nordström black hole[17] and the rotating Kerr-Newman black hole[18]. See also [19] for a different discussion. M. Angheben *et al* [20] also proposed a derivation of Hawking radiation by calculating the particles' classical action from the Hamilton-Jacobi equation, which is an extension of the complex path analysis of T. Padmanabhan *et al* [21]. Very recently, a new calculation concerning fermions' radiation from the stationary spherical symmetric black hole was done by R. Kerner and R. B. Mann in [22]. This method has been generalized to the more general and complicated spacetime background[23, 24, 25, 26, 27, 28, 29] and dynamical black hole[30].

In this paper, we will calculate the Fermions' Hawking radiation from the apparent horizon of FRW universe via tunneling formalism. Applying WKB approximation to the general covariant Dirac equation in FRW spacetime background, we also obtain the Hawking temperature of apparent horizon of FRW Universe. Our results obtained by taking the fermion tunneling into account support the arguments presented in the previous paper [1].

For convenience, we firstly give some results related to the FRW universe. The FRW metric is given by

$$ds^2 = -dt^2 + a^2(t) \left(\frac{dr^2}{1 - kr^2} + r^2(d\theta^2 + \sin^2\theta d\phi^2) \right) , \quad (1)$$

where $a(t)$ is the scale factor and $k = 1, 0$ and -1 represent the closed, flat and open universe respectively. Introducing $\tilde{r} = ar$, the FRW metric (1) can be rewritten as

$$ds^2 = h_{ab}dx^a dx^b + \tilde{r}^2 d\Omega_2^2 , \quad (2)$$

where $x^a = (t, r)$, $h_{ab} = \text{diag}(-1, \frac{a^2}{1-kr^2})$ and $d\Omega_2^2$ represents the line element of S^2 . The apparent horizon is defined by the equation

$$h^{ab}\partial_a\tilde{r}\partial_b\tilde{r} = 0 , \quad (3)$$

which gives us the location of apparent horizon explicitly as

$$\tilde{r}_A = \frac{1}{\sqrt{H^2 + k/a^2}} , \quad (4)$$

with $H = \dot{a}/a$ being the Hubble parameter. For simplicity, we will use the $(t, \tilde{r})$ coordinates, in which the FRW metric can be rewritten as

$$ds^2 = -\frac{1 - \tilde{r}^2/\tilde{r}_A^2}{1 - k\tilde{r}^2/a^2} dt^2 - \frac{2H\tilde{r}}{1 - k\tilde{r}^2/a^2} dt d\tilde{r} + \frac{1}{1 - k\tilde{r}^2/a^2} d\tilde{r}^2 + \tilde{r}^2 d\Omega_2^2 . \quad (5)$$

The inverse of the metric of the $(t, \tilde{r})$ parts is given by

$$g^{tt} = -1 \quad , \quad g^{t\tilde{r}} = -H\tilde{r} \quad , \quad g^{\tilde{r}\tilde{r}} = 1 - \tilde{r}^2/\tilde{r}_A^2 \quad . \quad (6)$$

Now we calculate the Fermions' Hawking radiation from the apparent horizon of the FRW universe via the tunneling formalism. Let us start with the massless spinor field Ψ obeyed the general covariant Dirac equation

$$-i\hbar\gamma^a e_a^\mu \nabla_\mu \Psi = 0 \quad , \quad (7)$$

where ∇_μ is the spinor covariant derivative defined by $\nabla_\mu = \partial_\mu + \frac{1}{4}\omega_\mu^{ab}\gamma_{[a}\gamma_{b]}$, and ω_μ^{ab} is the spin connection, which can be given in terms of the tetrad e_a^μ . The γ matrices are selected as

$$\begin{aligned} \gamma^0 &= \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix} \quad , \\ \gamma^1 &= \begin{pmatrix} 0 & \sigma^3 \\ \sigma^3 & 0 \end{pmatrix} \quad , \\ \gamma^2 &= \begin{pmatrix} 0 & \sigma^1 \\ \sigma^1 & 0 \end{pmatrix} \quad , \\ \gamma^3 &= \begin{pmatrix} 0 & \sigma^2 \\ \sigma^2 & 0 \end{pmatrix} \quad , \end{aligned}$$

where the matrices $\sigma^k (k = 1, 2, 3)$ are the Pauli matrices. According to the line element (5) and the inverse metric given in (6), the tetrad field can be selected to be

$$\begin{aligned} e_0^\mu &= (1, H\tilde{r}, 0, 0) \quad , \\ e_1^\mu &= (0, \sqrt{1 - k\tilde{r}^2/a^2}, 0, 0) \quad , \\ e_2^\mu &= (0, 0, \tilde{r}^{-1}, 0) \quad , \\ e_3^\mu &= (0, 0, 0, (\tilde{r}\sin\theta)^{-1}) \quad . \end{aligned}$$

Without loss of generality, we employ the following ansatz for spinor field with the spin up

$$\Psi = \begin{pmatrix} A(t, r, \theta, \phi) \\ 0 \\ B(t, r, \theta, \phi) \\ 0 \end{pmatrix} \exp\left[\frac{i}{\hbar}I(t, r, \theta, \phi)\right] \quad . \quad (8)$$

It should be noted that the spin-down case is just analogous. In order to apply the WKB approximation, we can insert the ansatz (8) for the spin-up spinor field Ψ into the general covariant Dirac equation (7). Dividing by the exponential term and neglecting the terms with $\hbar$, one can arrive at the following four equations

$$\begin{cases} iA(\partial_t I + H\tilde{r}\partial_{\tilde{r}}I) + B\sqrt{1 - k\tilde{r}^2/a^2}\partial_{\tilde{r}}I = 0 , \\ B\left(\frac{1}{\tilde{r}}\partial_{\theta}I + \frac{i}{\tilde{r}\sin\theta}\partial_{\phi}I\right) = 0 , \\ A\sqrt{1 - k\tilde{r}^2/a^2}\partial_{\tilde{r}}I - iB(\partial_t I + H\tilde{r}\partial_{\tilde{r}}I) = 0 , \\ A\left(\frac{1}{\tilde{r}}\partial_{\theta}I + \frac{i}{\tilde{r}\sin\theta}\partial_{\phi}I\right) = 0 . \end{cases} \quad (9)$$

Note that although A and B are not constant, their derivatives and the spin connections are all of the factor $\hbar$, so can be neglected to the lowest order in WKB approximation.

To carry out the separation of variables for the above equations, we must analysis the symmetries of the metric (5). For the metric (5), the Kodama vector is given by[1]

$$K^a = \sqrt{1 - k\tilde{r}^2/a^2} \left(\frac{\partial}{\partial t} \right)^a . \quad (10)$$

The Kodama vector in dynamical spacetime is of the same significance with the Killing vector in static spacetime. It should be noted that the Kodama vector is time-like, null and space-like as $\tilde{r} < \tilde{r}_A$, $\tilde{r} = \tilde{r}_A$ and $\tilde{r} > \tilde{r}_A$ respectively. Using the Kodama vector, one can define the energy measured by the Kodama observer

$$\omega = -K^a \partial_a I = -\sqrt{1 - k\tilde{r}^2/a^2} \partial_t I . \quad (11)$$

Using the definition of energy, the classical action can be separated as

$$I = - \int \frac{\omega}{\sqrt{1 - k\tilde{r}^2/a^2}} dt + R(\tilde{r}) + P(\theta, \phi) , \quad (12)$$

Substituting the above ansatz into Eq.(9) yields

$$\begin{cases} iA\left(-\frac{\omega}{\sqrt{1 - k\tilde{r}^2/a^2}} + H\tilde{r}\partial_{\tilde{r}}I\right) + B\sqrt{1 - k\tilde{r}^2/a^2}\partial_{\tilde{r}}I = 0 , \\ B\left(\frac{1}{\tilde{r}}\partial_{\theta}P(\theta, \phi) + \frac{i}{\tilde{r}\sin\theta}\partial_{\phi}P(\theta, \phi)\right) = 0 , \\ A\sqrt{1 - k\tilde{r}^2/a^2}\partial_{\tilde{r}}I - iB\left(-\frac{\omega}{\sqrt{1 - k\tilde{r}^2/a^2}} + H\tilde{r}\partial_{\tilde{r}}I\right) = 0 , \\ A\left(\frac{1}{\tilde{r}}\partial_{\theta}P(\theta, \phi) + \frac{i}{\tilde{r}\sin\theta}\partial_{\phi}P(\theta, \phi)\right) = 0 . \end{cases} \quad (13)$$

It is easy to see that P must be a complex function, which means it will yield a contribution to the imaginary part of the classical action. The contribution of P to the tunneling rate is

cancelled out when dividing the outgoing probability by the ingoing probability because P is completely the same for both the outgoing and ingoing solutions. It is no need to solve the equations about the complex function P . So it is essential to work out the imaginary part of $R(\tilde{r})$. From the first and third formulae of the above equation, there will be a non-trivial solution for A and B if and only if the determinant of the coefficient matrix vanishes, which results

$$\partial_{\tilde{r}} I = \frac{\omega}{(\tilde{r}^2/\tilde{r}_A^2 - 1)\sqrt{1 - k\tilde{r}^2/a^2}} \left(H\tilde{r} \pm \sqrt{1 - k\tilde{r}^2/a^2} \right) , \quad (14)$$

where the $+/-$ sign corresponds to the incoming/outgoing solutions respectively. It should be noted that the imaginary part of $R_{\pm}$ can be calculated using the above equation. Integrating the pole at the apparent horizon as in Refs.[31, 32], we have

$$\begin{aligned} \text{Im} R_+ &= \text{Im} \int \frac{\omega}{(\tilde{r}^2/\tilde{r}_A^2 - 1)\sqrt{1 - k\tilde{r}^2/a^2}} \left(H\tilde{r} + \sqrt{1 - k\tilde{r}^2/a^2} \right) d\tilde{r} \\ &= \pi\omega\tilde{r}_A , \\ \text{Im} R_- &= \text{Im} \int \frac{\omega}{(\tilde{r}^2/\tilde{r}_A^2 - 1)\sqrt{1 - k\tilde{r}^2/a^2}} \left(H\tilde{r} - \sqrt{1 - k\tilde{r}^2/a^2} \right) d\tilde{r} \\ &= 0 . \end{aligned} \quad (15)$$

In the WKB approximation, the tunnelling probability is related to the imaginary part of the action as

$$\Gamma = \frac{P_{\text{in}}}{P_{\text{out}}} = \frac{\exp[-2(\text{Im} R_+ + \text{Im} P)]}{\exp[-2(\text{Im} R_- + \text{Im} P)]} = \exp[-2\pi\omega\tilde{r}_A] . \quad (16)$$

Comparing the tunneling probability and the thermal spectrum $\Gamma = \exp[-\omega/T]$, Hawking temperature associated with the apparent horizon can be determined as

$$T = \frac{1}{2\pi\tilde{r}_A} . \quad (17)$$

The Hawking temperature associated with the apparent horizon of FRW universe obtained by using the fermions tunneling method takes the same form as that obtained in [1]. It should be noted that the Kodama observer is inside the apparent horizon. The results indicate that the Kodama observer does see a thermal spectrum with temperature $T = 1/2\pi\tilde{r}_A$, which is caused by fermions tunneling from the outside apparent horizon to the inside apparent horizon. The results in the present paper together with that in [1] fill in the gap existing in the literatures investigating the relationship between the first law of thermodynamics and Friedmann equations of FRW universe.

On the other hand, the higher terms about ω are neglected in our derivation and the expression (16) for tunneling probability implies the pure thermal radiation. When energy conservation is taken into account, the emission spectrum is no longer purely thermal and contains the higher terms about ω .

ACKNOWLEDGEMENT

This work was supported by the National Natural Science Foundation of China and Cuiying Project of Lanzhou University.

-
- [1] Rong-Gen Cai, Li-Ming Cao and Ya-Peng Hu, arXiv:0809.1554.
 - [2] T. Jacobson, Phys. Rev. Lett. **75**, 1260 (1995).
 - [3] C. Eling, R. Guedens and T. Jacobson, Phys. Rev. Lett. **96**, 121301 (2006).
 - [4] R. G. Cai and L. M. Cao, Phys. Rev. D **75**, 064008 (2007).
 - [5] R. G. Cai and S. P. Kim, JHEP **0502**, 050 (2005).
 - [6] M. Akbar and R. G. Cai, Phys. Rev. D **75**, 084003 (2007).
 - [7] M. Akbar, R. G. Cai, Phys.Lett. B **635**, 7 (2006).
 - [8] M. Akbar, R. G. Cai, Phys. Lett. B **648**, 243 (2007).
 - [9] R. G. Cai and L. M. Cao, Nucl. Phys. B **779**, 1 (2007).
 - [10] R. G. Cai, L. M. Cao and Y. P. Hu, JHEP **0808**, 090 (2008).
 - [11] X. H. Ge, Phys. Lett. B **651**, 49 (2007).
 - [12] Y. G. Gong, A. Z. Wang, Phys. Rev. Lett. **99**, 211301 (2007).
 - [13] S. F. Wu, B. Wang, G. H. Yang, Nucl. Phys. B **799**, 330 (2008).
 - [14] T. Zhu, J. R. Ren and S. F. Mo, arXiv:0805.1162.
 - [15] M. Akbar, arXiv:0808.0169.
 - [16] M.K.Parikh and F.Wilczek, Phys.Rev.Lett.**85**, 5042 (2000).
 - [17] J.Zhang and Z.Zhao, JHEP **0510**, 055 (2005).
 - [18] J.Zhang and Z.Zhao, Phys.Lett.B **618**, 14 (2005); J.Zhang and Z.Zhao, Phys.Lett.B **638**, 110 (2006).
 - [19] Q. Q. Jiang, S. Q. Wu, X. Cai, Phys.Rev. D **73**, 064003 (2006).

- [20] M.Angheben, M.Nadalini, L.Vanzo and S.zerbini, JHEP **0505**, 014 (2005).
- [21] K.Srinivasan and T.Padmanabhan, Phys.Rev.D **60**, 24007 (1999); S.Shankaranarayanan, T.Padmanabhan and K.Srinivasan, Class.Quant.Grav. **19**, 2671 (2002).
- [22] R.Kerner and R.B.Mann, Class.Quant.Grav. **25**, 095014 (2008).
- [23] R. Li, J. R. Ren, Phys. Lett. B, **661**, 370 (2008).
- [24] R. Li, J. R. Ren and S. W. Wei, Class. Quantum Grav. **25**, 125016 (2008).
- [25] R. Kerner, R.B. Mann, Phys. Lett. B, **665**, 277 (2008).
- [26] D. Y. Chen, Q. Q. Jiang, X. T. Zu, Phys. Lett. B, **665**, 106 (2008).
- [27] D. Y. Chen, Q. Q. Jiang, X. T. Zu, arXiv:0803.3248.
- [28] Q. Q. Jiang, Phys. Rev. D **78**, 044009 (2008).
- [29] Q. Q. Jiang, Phys. Lett. B, **666**, 517 (2008).
- [30] R. Di Criscienzo and L. Vanzo, Euro. Phys. Lett. **82**, 60001 (2008).
- [31] R. Kerner and R. B. Mann, Phys. Rev. D **73**, 104010 (2006).
- [32] P. Mitra, Phys. Lett. B **648**, 240 (2007).